



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/06	A1	(11) International Publication Number: WO 97/23601 (43) International Publication Date: 3 July 1997 (03.07.97)
<p>(21) International Application Number: PCT/US95/16843</p> <p>(22) International Filing Date: 21 December 1995 (21.12.95)</p> <p>(71) Applicant: GENZYME CORPORATION [US/US]; One Kendall Square, Cambridge, MA 02139 (US).</p> <p>(72) Inventors: WAN, Nick, C.; 11 Oakland Avenue, Newton, MA 02166 (US). McNEILLY, David, S.; 51 Birch Street, Saugus, MA 01906 (US). CHRISTOPHER, Charles, W.; 107 Main Street, Rockport, MA 01966 (US).</p> <p>(74) Agents: GOSZ, William, G. et al.; Genzyme Corporation, One Mountain Road, Framingham, MA 01701-9322 (US).</p>	<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: METHOD FOR LYSING CELLS</p> <p>(57) Abstract</p> <p>This invention relates to a method for lysing cells. The method comprises simultaneously flowing a cell suspension and a lysis solution through a static mixer, wherein the cells exit the static mixer lysed. In another aspect of the present invention, the invention relates to a method for precipitating cell components, protein, and nucleic acids from a cell lysate or other solution containing precipitable material. The method comprises simultaneously flowing a cell lysate or other protein containing solution and precipitating solution through a static mixer, wherein the lysate or protein solution exits the static mixer with its precipitable components precipitated. In another aspect of the present invention, the invention relates to a method where the two above-mentioned methods are combined by using static mixers in series.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR LYSING CELLS

5 Background of the Invention

Increasing attention has been focused on the delivery of DNAs as therapeutic agents (i.e., DNA gene therapy) for the treatment of genetic diseases and for genetic immunization.

10 Because of safety concerns with using potentially infectious viruses, researchers have been studying alternatives to viruses, using naked DNA or other non-viral methods of DNA delivery. One of the most promising non-viral methods in DNA therapy is the use of cationic lipids as delivery vehicles

15 (Felgner, *Proc. Natl. Acad. Sci.*, 84:7413-7417, 1987). The cationic lipids adsorb to negatively charged DNA and facilitate entry of the DNA into target cells. Successful delivery of genes via lipids into airway epithelia of rodents (Hyde, *Nature*, 362:250-255, 1993) have been reported. If future 'gene

20 therapy' proves to be effective, huge quantities of plasmids or appropriate DNA will be needed. However, current methods for isolating limited amounts of DNA may impede progress in this field.

Generally, the first step in making copies of a gene of

25 interest starts with inserting the appropriate portion of DNA into plasmids which are then replicated in "host cells". Host cells include eukaryotic and prokaryotic cells. Once a sufficient amount of plasmids are produced in the host cells, the host cells are lysed to release the internally replicated

30 plasmids.

Numerous mechanical methods for lysing cells have been developed and published. They include pressure, cavitation, sonic or ultrasonic waves, mechanical shaking with abrasives or grinding (Sadeva, *Biopharm* 7(2):26-36, 1994).

35 However, few of these methods are suitable for recovery of DNA or other shear sensitive materials from cells. Chemical

(Foster, *Biotechnol.* 10:273-277, 1992) and enzymatic (Andrews and Asenjo, *Tibtech.* 5:273-277, 1987) treatments, are more gentle and can be used for the recovery of shear sensitive products, such as DNA.

5 Generally, methods of plasmid isolation involve lysing the plasmid-containing cells with alkali (i.e. caustics) or enzymes in a test tube by gently inverting the tube several times. After the cells have been lysed, the cell lysate is mixed with a precipitating solution in a test tube by again gently
10 inverting the test tube several times. The object is to precipitate contaminating cell components, e.g. genomic DNA, proteins, and protein-nucleic acid complexes, leaving the plasmids in solution. The plasmids can then be separated from the precipitate by centrifugation. While this approach
15 can be used for small scale isolation, it is not practical for large scale production because the lysate is extremely viscous after the release of genomic DNA. Although, effective and gentle mixing of large quantities of viscous material can be achieved by using special agitators/mixers (e.g., planetary
20 mixers), these types of equipment are expensive and are not easily automated. In addition, the time of exposure of each plasmid to alkali in the large volume of solution is difficult to control and can vary a great deal and this may affect plasmid quality. Furthermore, it is crucial to handle the lysate very
25 gently because any sheared genomic DNA may contaminate the sample and render later plasmid purification extremely difficult.

 A need exists to develop an effective, economical, and automatable method for isolating high quality plasmids on a
30 large scale to meet the future demand.

Summary of the Invention

 This invention relates to the use of static mixers in a
35 novel manner, such as to rapidly lyse large amounts of cells. The lysis method comprises simultaneously flowing a

cell suspension and a lysis solution through a static mixer, wherein the cells exit the static mixer lysed. A key advantage of the present invention is that multi-liter amounts of solution containing multi-gram amounts of cells
5 can be lysed rapidly, making large scale biological procedures involving cell lysis feasible.

In another aspect of the present invention, the invention relates to a rapid method for precipitating cell components, protein, and nucleic acids from a cell lysate or
10 other solution to be precipitated, using a static mixer. The method comprises simultaneously flowing a cell lysate or other protein-containing solution and a precipitating solution through a static mixer, wherein the lysate or protein solution exits the static mixer with its precipitable
15 components precipitated.

In another aspect of the present invention, the invention relates to a method where the two above-mentioned methods are combined by using static mixers in series, i.e., by first lysing the cells through a first static
20 mixer and then precipitating the cell lysate through a second static mixer.

In another aspect of the present invention, the invention relates to a method for releasing intact usable plasmids from plasmid-containing cells, comprising
25 simultaneously flowing a solution containing the plasmid-containing cells and a lysate solution through a static mixer wherein the plasmid-containing cells exit the static mixer lysed and the plasmids released. In addition, the method of releasing a plasmid further includes a preliminary step,
30 prior to the lysing step, of simultaneously diluting the cells to be lysed to an optimal cell density by flowing the plasmid-containing cell suspension and a solution through a static mixer prior to simultaneously flowing the diluted cell suspension with a lysis solution through another static
35 mixer (i.e., a second static mixer) with the same results. Furthermore, this method can include the precipitation

step, explained above, where the cell lysate of the second static mixer is combined with a precipitating solution through a third static mixer.

5

Brief Description of the Drawings

Figure 1 is an illustration of the method of the present invention where cells are passed through a static mixer simultaneously with a lysis solution, the result being that the cells exit the static mixer lysed.

10

Figure 2 is an illustration of the method of the present invention where a cell lysate (or protein solution) is passed through a static mixer simultaneously with a precipitating solution the result being that the cell lysate (or protein solution) exits the static mixer precipitated.

15

Figure 3 is an illustration of the method of the present invention where cells are first passed through a static mixer simultaneously with a cell diluting solution, then the mixture exiting the first static mixer is passed through a second static mixer along with a lysis solution, then that mixture is passed through a third static mixer simultaneously with a precipitating solution.

20

Detailed Description of the Invention

25

This invention is based upon the discovery that static mixers could be used to lyse cells containing plasmids, releasing the plasmids from the cells. The advantage of using such a device is that large volumes of cells can be gently and continuously lysed in-line using the static mixer and that other static mixers could be placed in-line to accomplish other functions such as dilution and precipitation. This method greatly simplifies the process of isolating plasmids from large volumes of material such that plasmid DNA is not damaged by the process. Previous methods of plasmid isolation involving caustic lysing and precipitation, which involved expensive and

30

35

specialized equipment, were not practical for large scale plasmid purification.

5 The method of the present invention can be used to lyse any type of cell (i.e., prokaryotic or eukaryotic) for any purpose related to lysing, such as releasing desired nucleic acids or proteins from target cells to be subsequently purified. In a preferred embodiment, the method of the present invention is used to lyse host cells containing plasmids to release plasmids.

10 The term "lysing" refers to the action of rupturing the cell wall and/or cell membrane of a cell which is in a buffered solution (i.e., cell suspension) through chemical treatment using a solution containing a lysing agent. Lysing agents include for example, alkali, detergents,
15 organic solvents, and enzymes. In a preferred embodiment, the lysis of cells is done to release intact plasmids from host cells. For purposes of the present invention the term "simultaneously" referring to the passage of substances through a static mixer, means that
20 the subject substances are passing through the static mixer at approximately the same time.

For purposes of the present invention the term "flowing" refers to the passing of a liquid at a particular flow rate (e.g., liters per minute) through the static mixer,
25 usually by the action of a pump. It should be noted that the flow rate through the static mixer is believed to affect the efficiency of lysis, precipitation and mixing.

Suitable static mixers useful in the method of the present invention include any flow through device referred
30 to in the art as a static or motionless mixer of a length sufficient to allow the processes of the present invention. For example, for the purpose of lysing cells, the static mixer would need to have a length which would provide enough contact time between the lysing solution and the cells to
35 cause the lysis of the subject cells during passage through the mixer. In a preferred embodiment, suitable static

Escherichia coli (*E. coli*) cells that had been grown in a nutrient medium to a high cell density, the culture was then diluted directly by flowing the cells through a 1/2" x 27", 32-element, Kenics static mixer (purchased from
5 Chemineer, N. Andover MA) along with a resuspending solution (50mM Tris/HCl, 10mM EDTA, 100 mg RNaseA/ml, pH 8.0) until the exiting cells had an optical density of 25-40 at 600nm, as read by a spectrophotometer.

10 The diluted cell suspension exiting the first static mixture was then flowed through a second static mixer (same type of mixer as above) along with a lysis solution (200mM NaOH, 1% weight/volume SDS,) at a rate of 100-1,300 ml/min. The cells exited the second static mixer
15 lysed (i.e., cell lysate).

The cell lysate exiting the second static mixer was then flowed through a third static mixer (same type as above) with a precipitating solution (2.6 M potassium acetate, pH 5.2) at a similar flow rate as passed through
20 the second static mixer. The exiting suspension contained precipitated *E. coli* genomic DNA, proteins, and insoluble debris and the soluble plasmids. About 45 liters of mixture was collected from the third static mixer and processed. Approximately 1 gram of plasmid was isolated
25 from this run.

All of the static mixers above were connected in series by tubing and the respective solutions were flowed or pumped through the static mixers by peristaltic pumps (Cole-Parmer, Chicago IL).
30

THE INVENTION CLAIMED IS:

1. A method of lysing cells, comprising simultaneously
5 flowing a cell suspension and a lysis solution through
a static mixer, wherein the cells exit the static mixer
lysed.
2. The method of claim 1 wherein the lysis solution is a
10 solution containing a lysis agent selected from the
group consisting of an alkali, a detergent, an organic
solvent, and an enzyme or a mixture thereof.
3. The method of claim 1 wherein the cells are plasmid-
15 containing cells.
4. A method of precipitating cellular components from a
solution, comprising simultaneously flowing a cell
20 lysate and a precipitating solution through a static
mixer, wherein the cellular components exit the static
mixer precipitated.
5. The method of claim 4 wherein the precipitating
25 solution is a solution containing a precipitating agent
selected from the group consisting of sodium dodecyl
sulfate and potassium acetate or a mixture thereof.
6. The method of claim 4 wherein the cell lysate
contains plasmids.
- 30 7. A method of releasing plasmids from plasmid-
containing cells, comprising simultaneously flowing a
suspension containing the plasmid-containing cells
and a lysis solution through a static mixer wherein
35 the plasmid-containing cells exit the static mixer
lysed and plasmids released from the cells.

8. The method of claim 7 wherein the lysis solution is a solution containing a lysis agent selected from the group consisting of an alkali, a detergent, an organic solvent, and an enzyme or mixture thereof.
- 5
9. A method of releasing plasmids from plasmid-containing cells, comprising:
- 10
- a. simultaneously flowing a suspension containing the plasmid-containing cells and a buffered solution through a first static mixer; and
- b. simultaneously flowing the product of step a) and a lysis solution through a second static mixer wherein the plasmid-containing cells exit the static mixer lysed and plasmids released
- 15
- from the cells.
10. The method of claim 9 wherein the lysis solution is a solution containing a lysis agent selected from the group consisting of an alkali, a detergent, an organic solvent, and an enzyme or mixture thereof.
- 20
11. A method of releasing plasmids from plasmid-containing cells, comprising:
- 25
- a. simultaneously flowing a suspension containing the plasmid-containing cells and a buffered solution through a first static mixer;
- b. simultaneously flowing the product of step a) and a lysis solution through a second static mixer wherein the plasmid-containing cells exit the static mixer lysed and the plasmid released
- 30
- into the lysate; and
- c. simultaneously flowing the product of step b) through a third static mixer with a precipitating solution wherein the cellular components exit the static mixer precipitated, leaving the
- 35
- plasmid in solution.

12. The method of claim 11 wherein the lysis solution is
a solution containing a lysis agent selected from the
group consisting of an alkali, a detergent, an organic
solvent, and an enzyme or mixture thereof.
13. The method of claim 11 wherein the precipitating
solution is a solution containing a precipitating agent
selected from the group consisting of sodium dodecyl
sulfate and potassium acetate or mixture thereof.
14. A method of releasing plasmids from plasmid-
containing cells, comprising:
- a. simultaneously flowing a suspension containing
the plasmid-containing cells and a lysis solution
through a first static mixer wherein the
plasmid-containing cells exit the static mixer
lysed and the plasmid released; and
- b. simultaneously flowing the product of step a)
through a second static mixer with a
precipitating solution wherein the product of
step a) exits the static mixer precipitated and
the plasmid soluble.
15. The method of claim 14 wherein the lysis solution is
a solution containing a lysis agent selected from the
group consisting of an alkali, a detergent, an organic
solvent, and an enzyme or mixture thereof.
16. The method of claim 14 wherein the precipitating
solution is a solution containing a precipitating agent
selected from the group consisting of sodium dodecyl
sulfate and potassium acetate or mixture thereof.

FIGURE 1




FIGURE 2





FIGURE 3



INTERNATIONAL SEARCH REPORT

later. national application No.
PCT/US95/16843

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C12N 1/06

US CL :435/259, 320.1, 287

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/259, 320.1, 287

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
KIRK-OTMHER

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE DIALOG, MEDLINE, WPIDS

search terms: lysis, lysing, static mixer, plasmid?, recover?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	Journal of Chemical Technology and Biotechnology, Volume 54, issued 1992, Papamichael et al., "Continuous Aqueous Phase Extraction of Proteins: Automated Processing and Recycling of Process Chemicals", pages 47-55, especially page 48.	1-6 ----- 7-16
X --- Y	US, A, 4,462,940 (HANISCH et al.) 31 July 1984, column 10, lines 20-38.	1-3, 5-8 ----- 4-6, 10-16
Y	US, A, 4,900,677 (HEWITT) 13 February 1990, column 9, lines 21-60.	1-16
Y	US, A, 5,208,160 (KIKYOTANI et al.) 04 May 1993, Example 5 and comparative Example.	1-16

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

11 APRIL 1996

Date of mailing of the international search report

29 APR 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

IRENE MARX

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/16843

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,096,818 (DEBONVILLE) 17 March 1992, Example 1.	1-16
Y	US, A, 4,997,932 (REARDON et al.) 05 March 1991. See whole document, in particular, column 6, lines 5-19.	1-16
X	US, A, 4,450,103 (KONRAD et al.) 22 May 1984. See entire document, especially Example 3.	1-3, 7-10
—		
Y		4-6, 11-16
Y	Enzyme and Microbial Technology, Volume 6, number 7, issued 06 July 1984, Veide et al., "Continuous extraction of beta-D-galactosidase from Escherichia coli in an aqueous two-phase system: effects of biomass concentration on partitioning and mass transfer, pages 325-330, especially page 326.	1-16
Y	RODRIGUEZ ET AL., 'RECOMBINANT DNA TECHNIQUES: AN INTRODUCTION" published 1983 by The Benjamin/Cummings Publishing Company, Inc., (CA), see pages 37-46, especially, pages 45-46.	1-16

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.